

Cluster Analysis for Percolation on Two Dimensional Fully Frustrated System

Giancarlo Franzese

Dipartimento di Scienze Fisiche, Università di Napoli, Mostra d'Oltremare Pad.19 I-80125 Napoli

Italy

and INFN - unità di Napoli

(February 1, 2008)

Abstract

The percolation of Kandel, Ben-Av and Domany clusters for 2d fully frustrated Ising model is extensively studied through numerical simulations. Critical exponents, cluster distribution and fractal dimension of percolative cluster are given.

I. INTRODUCTION

A 2d fully frustrated (FF) Ising model is a model with Ising spins ± 1 where the interactions between nearest neighbor spins have modulus $J > 0$ and sign ± 1 (ferro / antiferromagnetic interactions) and where the signs are chosen in such a way that every *plaquette* (i.e. the elementary cell of square lattice) is *frustrated*, i.e. every plaquette has an odd number of -1 interactions so that the four spins of the plaquette cannot simultaneously satisfy all four interactions. In Fig.1.1 we give an example of such a deterministic interaction configuration. In a plaquette of the FF model we can have only one or three satisfied interactions. The FF model has an analytical solution [1] and a critical temperature at $T_c = 0$.

Since single-spin dynamics for FF suffers critical slowing down, a fast cluster dynamics was introduced by Kandel, Ben-Av and Domany (KBD) in Ref. [2].

The KBD-clusters are defined by choosing stochastically on each plaquette of a checker-board partition of a square lattice one bond configuration between the three shown in Fig.1.2.

The probability of choice depends on spins configuration on the plaquette and it is a function of temperature (*correlated site-bond percolation* [3]). When there is only one satisfied interaction the zero-bond configuration is chosen with probability one. When three interactions are satisfied the zero-bond configuration is chosen with probability $P_0 = e^{-4J/(kT)}$ (where k is the Boltzmann constant and T the absolute temperature), the bond configuration with two parallel bonds on two satisfied interactions is chosen with probability $P_1 = 1 - P_0$ and the third bond configuration has zero probability. Two sites are in the same cluster if they are connected by bonds. For sake of simplicity from now on we choice $J/(kT) = 1/T$.

Ref. [2] has stimulated several works [4,5] that pay attention mainly to dynamics and to number of clusters and clusters sizes. In [5] numerical simulations on relatively large FF lattice sizes (number of sites $N = 60^2 \div 120^2$) supported the idea that the KBD-clusters represent spin correlated regions (like Coniglio-Klein clusters [6] do in Ising model) and consequently percolation temperature T_p coincides with critical temperature T_c , percolation exponents coincide with critical ones and KBD-clusters at T_p are 2d self-avoiding walks

(SAW) at θ point. [7]

In this paper we extensively study percolative features of KBD-clusters, considering very large lattice sizes ($N = 100^2 \div 400^2$), and give numerical results on critical exponents, cluster distribution and fractal dimension at percolation point.

II. CRITICAL EXPONENTS AND PERCOLATION POINT

We consider finite systems with increasing size ($L = 100 \div 400$) with periodic boundary conditions.

A cluster percolates when it connects two opposed system sides. For every size L there is a percolation temperature $T_p(L)$. With T_p (without any argument) we mean the percolation temperature in the thermodynamic limit i.e. $T_p(L) \rightarrow T_p$ for $L \rightarrow \infty$. In this limit percolating clusters are present at $T \leq T_p$ but not at $T > T_p$. In Fig.2 we show typical clusters at several temperatures for a finite system with size $L = 60$.

For every L we have studied the mean cluster size $S = \sum_s s^2 n_s$ (where s is the cluster size, n_s the number of cluster of size s per lattice site and the sum is extended over all finite clusters), the percolation probability $P = 1 - \sum_s s n_s$, the number of cluster $N_c = \sum_s n_s$, the number of bonds per lattice site N_b , the mean size of the largest (percolating) cluster S_I and the mean size of the second largest (percolating) cluster S_{II} . These quantities are shown in Fig.3 for $L = 100 \div 400$. Let's note that for $T \rightarrow 0$ the bonds cover 50% of lattice interactions (that is the random-bond percolation threshold on square lattice), S_{II} goes to a finite value (like predicted by KBD [2] and already verified in Ref. [4]) and occupies almost 35% of the lattice, and that S_I occupies almost 65% of the lattice. At $T = 0$ only two clusters survive, as shown in Fig.2 e) and f).

Now we will give numerical estimates of critical exponents that characterize the KBD-cluster percolation.

We know [8] that in the thermodynamic limit the mean cluster size diverges for $T \rightarrow T_p$, the percolation probability goes to zero in the limit $T \rightarrow T_p^-$ and the number of cluster goes

to zero for $T \rightarrow T_p^+$.

We assume that near T_p the *connectivity length* ξ (i.e. the typical linear cluster size) diverges like $\xi \sim |e^{-2/T} - e^{-2/T_p}|^{-\nu}$, the mean cluster size diverges like $S \sim |e^{-2/T} - e^{-2/T_p}|^{-\gamma}$, the percolation probability goes to zero like $P \sim |e^{-2/T} - e^{-2/T_p}|^\beta$ and the number of cluster goes to zero like $N_c \sim |e^{-2/T} - e^{-2/T_p}|^{2-\alpha}$. The last relations are definitions of critical exponents α , β , γ and ν .

By standard finite-size scaling considerations [8] we can make the *ansatz*

$$S \sim L^{\gamma/\nu} f_S(|e^{-2/T} - e^{-2/T_p}| L^{1/\nu}) \quad (1)$$

$$P \sim L^{-\beta/\nu} f_P(|e^{-2/T} - e^{-2/T_p}| L^{1/\nu}) \quad (2)$$

and

$$N_c \sim L^{(\alpha-2)/\nu} f_{N_c}(|e^{-2/T} - e^{-2/T_p}| L^{1/\nu}) \quad (3)$$

where $f_S(x)$, $f_P(x)$ and $f_{N_c}(x)$ are *universal functions*, i.e. independent by L .

Via data-collapse (see Fig.4) we estimate the parameters $e^{-2/T_p} = 0.0000$, $\alpha = 0.1$, $\beta = 0.00$, $\gamma = 2.00$, and $\nu = 1.00$ with error of one unit in the last given digit. Therefore the scaling relation $\alpha + 2\beta + \gamma = 2$ and the hyperscaling relation $2 - \alpha = \nu d$ are satisfied with good approximation.

In Tab.1 we give numerical estimates of $T_p(L)$. The data are obtained taking for $L = 100, 200, 300, 400$ the values of $T_p(L)$ at which the S data in a log-log plot vs. $|e^{-2/T} - e^{-2/T_p(L)}|$ follow two parallel straight lines (one above and one below $T_p(L)$) with slopes in good agreement with $\gamma = 2$ and then best-fitting these values as $e^{-2/T_p(L)} \sim 1/L$.

III. FRACTAL DIMENSION AND CLUSTER DISTRIBUTION

Let's now consider the fractal dimension D of the percolating cluster. From the scaling invariance hypothesis [8] we know that $P \sim \xi^{D-d}$, then we obtain $D = d - \beta/\nu$ (hyperscaling).

In present case we have $\beta = 0$, then $D = d = 2$. The same result is obtained from the scaling relation $\beta + \gamma = D\nu$.

This is confirmed by the analysis of cluster distribution (see Fig.5). The scaling invariance hypothesis [8,9] gives for $T \rightarrow T_p$ and $s \rightarrow \infty$

$$n_s = s^{-\tau} f_{n_s}(|e^{-2/T} - e^{-2/T_p}| s^\sigma) \quad (4)$$

with $\tau = 1 + d/D$, $\sigma = 1/(\nu D)$ and $f_{n_s}(x)$ universal function.

From data-collapse for n_s near T_p (see Fig.6.a) we obtain numerical estimates of parameters. The data in Fig.6.a are chosen in such a way that the quantity $(e^{-2/T} - e^{-2/T_p})L^{1/\nu}$ (with $e^{-2/T_p} = 0$ and $\nu = 1$) is a constant with $T \simeq T_p(L)$ for every considered L . The results are $\tau = 2.00$ and $\sigma = 0.50$ (with error of one unit in the last digit), that, with the definitions of τ and σ , give $D = 2$ and $\nu = 1$. On the other hand these values of τ and σ satisfies the relation $\sigma(2 - \alpha) = \tau - 1$, $\sigma\beta = \tau - 2$, $\sigma\gamma = 3 - \tau$. [8]

From Fig.6.a we see that the universal function $f_{n_s}(x)$ is a bell-shaped curve for $T \simeq T_p(L)$. For temperatures slightly below $T_p(L)$ (Fig.6.b) $f_{n_s}(x)$ is shifted, while for temperatures slightly above $T_p(L)$ (Fig.6.c) $f_{n_s}(x)$ changes dramatically its shape.

Away from $T_p(L)$ we know [8] that is valid the relation

$$\log n_s \sim -s^\zeta \quad (5)$$

for $s \rightarrow \infty$, with $\zeta = 1$ above $T_p(L)$ and $\zeta = 1 - 1/d = 1/2$ below $T_p(L)$. This relation is confirmed with reasonable approximation by our numerical simulations, as shown in Fig.7. Let's note that, while the exponent $\zeta = 1$ above $T_p(L)$ is good for a wide range of s ($s = 2000 \div 8000$ for $L = 100$), the exponent $\zeta = 1/2$ below $T_p(L)$ is good for a smaller s range ($s = 2000 \div 4400$ for $L = 100$) since finite-size effect become more important below $T_p(L)$. The smaller T , the smaller s range is.

A direct way to estimate the fractal dimension D is given through its definition

$$s \sim R^D \quad (6)$$

for $T = T_p$, with R radius of gyration of the cluster of size s . We know [8] that cluster dimension deviates from D away from T_p , becoming the Euclidean dimension d below T_p and a value smaller than D above T_p . This is true because eq.(6) is valid within the connectivity length ξ for all temperatures, but ξ goes to zero away from T_p . Unfortunately data about this relation are difficult to analyze. Indeed near $T_p(L)$ for every finite system with $L \leq 120$ it seems that D is almost $7/4 = 1.75$ (the fractal dimension of a SAW at θ point), but for larger L (see Fig.8 and Tab.1) the fractal dimension D grows slowly to the asymptotic value 2.

IV. CONCLUSIONS

We have numerically investigated the KBD-cluster percolation problem in 2d FF Ising model. From our simulation we found that, within numerical errors, this correlated site-bond percolation satisfies scaling and hyperscaling relations and have, in the thermodynamical limit, a percolation temperature $T_p = 0$ and the exponents $\alpha = 0$, $\beta = 0$, $\gamma = 2$, $\nu = 1$, $\tau = 2$, $\sigma = 1/2$, $\zeta(T > T_p(L)) = 1$, $\zeta(T < T_p(L)) = 1/2$. Moreover at T_p clusters are compact (fractal dimension $D = 2$). Therefore now we can correct the conclusion of Ref. [5] and say that, since $T_p = T_c$ and ν is equal to spin correlation exponent [10], the site connectivity length ξ goes like the spin correlation length diverging at zero temperature. Although the exponent γ is different the coincidence between ξ and correlation length is enough to give an efficient Monte Carlo cluster dynamics. [11]

ACKNOWLEDGMENTS

The author is indebted to Antonio Coniglio and to Vittorio Cataudella for many illuminating discussions and a careful reading of the manuscript.

The computation have been done on DECstation 3000/500 with Alpha processor and DECsystem 5000/200 with RISC processor.

REFERENCES

- [1] J.Villain, J. Phys. C **10**, 1717 (1977);
G.Forghas, Phys. Rev. B **22**, 4473 (1980).
- [2] D.Kandel, R.Ben-Av and E.Domany, Phys. Rev. Lett. **65**, 941 (1990);
D.Kandel and E.Domany, Phys. Rev. B **43**, 8539 (1991).
- [3] A.Coniglio, H.E.Stanley and W.Klein, Phys. Rev. Lett. **42**, 518 (1979).
- [4] W.Kerler and P.Rehberg, Phys. Rev. B **49**, 9688 (1994);
P.D.Coddington and L. Han, Phys. Rev. B **50**, 3058 (1994);
- [5] V.Cataudella, G.Franzese, M.Nicodemi, A.Scala and A.Coniglio, Phys. Rev. Lett. **47**,
381 (1994) and
Il Nuovo Cimento **16D** N.8, 1259 (1994).
- [6] A.Coniglio and W.Klein, J. Phys. A **13**, 2775 (1980).
- [7] A.Coniglio, N.Jan,I.Maijd and H.E.Stanley, Phys. Rev. B **35**, 3617 (1987);
B.Duplantier and H.Saleur, Phys. Rev. Lett. **59**, 538 (1987).
- [8] D.Stauffer, *Introduction to Percolation Theory* (Taylor & Francis, London, 1985).
- [9] M.D'Onorio De Meo, D.W.Heermann and K.Binder, J. Stat. Phys., **60**, 585 (1990).
- [10] as for clusters of parallel spin in 2d Ising model (Ref. [6]).
- [11] V.Cataudella, G.Franzese, M.Nicodemi, A.Scala and A.Coniglio, in print on Phys. Rev.
E (cond-mat/9604169)


TABLES

TABLE I. Numerical estimates of $T_p(L)$ and $D(L)$ for $L = 60 \div 400$. The way used to evaluate $T_p(L)$ and $D(L)$ give us confidence only on digit not in parentheses.

L	60	80	100	120	200	300	400
$T_p(L)$	0.51(7)	0.48(1)	0.45(6)	0.43(7)	0.39(2)	0.36(1)	0.34(2)
$D(L)$	1.7(2)	1.7(5)	1.7(7)	1.7(9)	1.8(2)	1.8(5)	1.8(6)

FIGURES

1)



2)




FIG. 1. 1) Example of 2d FF lattice: the spin are on the vertices; solid lines represent ferromagnetic interactions ($+J$) and dashed lines antiferromagnetic interactions ($-J$). 2) Plaquette bond configurations: a) zero bond; b) two parallel vertical bonds; c) two parallel horizontal bonds.




FIG. 2. Typical KBD-clusters on a FF lattice with size $L = 60$ with periodic boundary conditions: a) at $T = 1$; b) at $T = 0.65$; c) at $T = 0.53$ slightly above $T_p(L) \simeq 0.52$; d) at $T = 0.52 \simeq T_p(L)$; e) at $T \simeq 0$ (largest cluster); f) at $T \simeq 0$ (the second cluster).




FIG. 3. Mean cluster size S , percolation probability P , number of cluster N_c , number of bonds per lattice site N_b , mean size of largest cluster S_I and mean size of second largest cluster S_{II} vs. temperature T for lattice sizes $L = 100, 200, 300, 400$. The error bars (often included in symbols) are the statistical errors.




FIG. 4. Scaling for S , P and N_c following assumptions (1), (2) and (3) for data of systems with sizes $L = 100, 200, 300, 400$. The parameters $e^{-2/T_p} = 0.0000 \pm 0.0001$, $\alpha = 0.1 \pm 0.1$, $\beta = 0.00 \pm 0.01$, $\gamma = 2.00 \pm 0.01$ and $\nu = 1.00 \pm 0.01$ are such that the data for different sizes L collapse on single curves (one for each graph). These curves are, respectively, the universal functions f_S , f_P and f_{N_c} . The errors are estimated observing the range of parameters within which the data points collapse roughly well.




FIG. 5. Cluster distribution n_s vs. cluster size s for a system with size $L = 100$ at several temperatures T . Every bin is large 400 unities in cluster size. The percolation temperature for this system size is $T_p(L = 100) \simeq 0.46$ then for all $T \geq 0.47$, even if the highest bin is not empty, there are only no percolating clusters. Let's note that for all T above $T_p(L)$ it is $n_s \sim e^{-s}$ and that the distribution become symmetric for $T \rightarrow 0$.




FIG. 6. a) Scaling for n_s following assumption (4) with parameter $\tau = 2.00 \pm 0.01$ and $\sigma = 0.50 \pm 0.01$ for data of systems with sizes $L = 100, 200, 300, 400$. Each set of data is chosen at a temperature near the corresponding $T_p(L)$. As consequence for each temperature the quantity $(e^{-2/kT} - e^{-2/T_p})L^{1/\nu}$ with $T_p = 0$ and $\nu = 1$ is equal to 2.084. Every point in the graph is an average over 500 consecutive values of s for $L = 100$, 2000 for $L = 200$, 4500 for $L = 300$, 8000 for $L = 400$. b) As in part a) but for $(e^{-2/T} - e^{-2/T_p})L^{1/\nu} = 0.823$ (below $T_p(L)$). c) As in part a) but for $(e^{-2/T} - e^{-2/T_p})L^{1/\nu} = 3.567$ (above $T_p(L)$).




FIG. 7. Fit of $\log(-\log n_s L^2)$ vs. $\log(s/400)$ for a system with $L = 100$ and $T_p(L) \simeq 0.46$: a) at $T \simeq 0.47 > T_p(L)$ the slope is $\zeta \simeq 1$; b) at $T \simeq 0.45 < T_p(L)$ the slope is $\zeta \simeq 1/2$.




FIG. 8. Estimate of D from definition (6) vs. $1/L$ (see Tab.1): at $T = 0.517$ for $L = 60$, at $T = 0.481$ for $L = 80$, at $T = 0.450$ for $L = 100$, at $T = 0.437$ for $L = 120$, at $T = 0.389$ for $L = 200$, at $T = 0.360$ for $L = 300$ and at $T = 0.343$ for $L = 400$. The error bars are probably underestimated. The arrow heads for the asymptotic value of D .